# Numerical Approximation Of Two-Dimensional Stochastic Neural Field Equations With Finite Transmission Speed

## <u>Pedro Lima<sup>1</sup></u> and Tiago Sequeira<sup>1</sup>

<sup>1</sup> Department of Mathematics and CEMAT, Instituto Superior Técnico, Lisboa, Portugal

 $Presenting \ author: \ plima@math.tecnico.ulisboa.pt$ 

### Talk Abstract

Neural field equations are intended to model the synaptic interactions between neurons in a continuous neural network, called a neural field [1], [2]. This kind of integro-differential equations proved to be a useful tool for the spatiotemporal modeling of the neuronal activity from a macroscopic point of view, allowing the study of a wide variety of neurobiological phenomena, such as the processing of sensory stimuli. The aim of the present talk is to study the effects of additive noise in one- and two-dimensional neural fields, while taking into account finite signal transmission speed. A Galerkin-type method to approximate such models is presented, which applies the Fast Fourier Transformation to optimise the computational effort required to solve this type of equations. Numerical simulations obtained by this algorithm are presented and discussed.

**Keywords:** Neural field equation, stochastic differential equation, Galerkin method.

#### Acknowledgements

This work was supported by FCT, within project UIDB/04621/2020 (Center for Computational and Stochastic Mathematics https://doi.org/10.54499/U-IDB/04621/2020).

### References

- Amari S.L., Dynamics of pattern formation in lateral-inhibition type neural fields, Biol. Cybernet., 27 (2), 1977, P. 77–87.
- [2] Wilson, H.R. and Cowan, J.D., Excitatory and inhibitory interactions in localized populations of model neurons, Bipophys. J., 12, 1972, P. 1–24.